



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2015

---

## **Surgical Relevance of Corona Mortis and Clinical Outcome in Pelvic Trauma**

Jensen, K O ; Sprengel, K ; Mica, L ; Somlyay, L ; Jentzsch, T ; Werner, C M L

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-118045>

Journal Article


Published Version

Originally published at:

Jensen, K O; Sprengel, K; Mica, L; Somlyay, L; Jentzsch, T; Werner, C M L (2015). Surgical Relevance of Corona Mortis and Clinical Outcome in Pelvic Trauma. *Austin Journal of Anatomy:online*.

Anatomy-v2-id1033
Top
Title
Abstract
Abbreviations
Introduction
Patients and Methods
Results
Discussion
Conclusion
References

Justin  
Journal of  
Anatomy



Open Access

# Surgical Relevance of Corona Mortis and Clinical Outcome in Pelvic Trauma

Kai Oliver Jensen<sup>\*,#</sup>, Kai Sprengel<sup>#</sup>, Ladislav Mica, Lukas Somlyay, Thorsten Jentzsch and Clément ML Werner

Division of Trauma Surgery, University of Zurich, Switzerland <sup>#</sup>Both authors contribute equally

**\*Corresponding author:** Kai Oliver Jensen, Division of Trauma Surgery, University Hospital Zurich, University of Zurich, Switzerland, Raemistrasse 100, 8091 Zürich

**Received:** December 21, 2014; **Accepted:** March 17, 2015 **Published:** March 18, 2015

## Abstract

**Introduction:** The Corona Mortis (CM), also termed the crown of death, is an anastomosis between the obturator and the external iliac or inferior epigastric vessels. It has been accused to be a source of major bleeding complications in many instances. Not only pelvic surgeons know that dislocated fractures or iatrogenic injury may cause life-threatening bleeding. For the investigation of the incidence and surgical relevance of the CM, we conducted a retrospective study considering the intra- and peri-operative findings at our institution.

**Methods:** All consecutive patients, who underwent pelvic surgery due to an acetabular fracture or pelvic ring instability were analysed. Patients were excluded if an approach without standard visualization of the CM was used. The incidence of the CM and its associated bleeding complications and mortality was investigated as documented by the operative report and patients' medical records. After performing a two-tailed Kolmogorov-Smirnov test for normal distribution, a Mann-Whitney U test was used due to a nonparametric distribution to test nominal variables. P-values of <0.05 were considered statistically significant.

**Results:** We evaluated the data of 130 consecutive patients. The incidence of the CM was 41.5% (n=54). Neither a significant correlation between the presence of the CM and bleeding complications ( $p=0.068$ ), nor the mortality ( $p=0.068$ ) was observed. The mortality was also not significantly affected by the presence of the CM or bleeding complications ( $p=0.338$  and  $p=0.113$ , respectively).

**Conclusion:** This study points out the high incidence of the CM. However, it is not associated with bleeding complications or mortality. Therefore, it does not seem to be the main source of bleeding in pelvic trauma in case of delayed surgery and can be regarded as a controllable anatomic structure.

**Keywords:** Corona mortis; Pelvic fracture; Incidence; Bleeding; Complications; Mortality

## Abbreviations

CM: Corona Mortis; CT: Computed Tomography

## Introduction

The Corona Mortis (CM) is known as an anastomosis between the obturator and the external iliac or the inferior epigastric vessels. It is described as an anatomical variant with variable size (>1 mm to <4 mm), distance to the ramus pubis (40 mm to 96 mm) and incidence (25% to 83 %) [1-4]. During pelvic surgeries, most surgeons use an anterior approach for the reduction and fixation of fractures of the pelvic ring and acetabulum [5-8]. On the one hand, the CM is a cause of major bleeding complications due to dislocated fractures and, on the other hand, it is a potential risk factor for significant haemorrhage by iatrogenic, intraoperative lesions, which may be limb- and lifethreatening [3,9-12]. However, the current literature is based on cadaver, angiographic and Computed Tomography (CT) studies. To determine the incidence and clinical relevance of the CM, we conducted a retrospective study based on intra- and peri-operative findings at our institution.

Received: December 21, 2014; Accepted: March 17, 2015; Published: March 18, 2015

File failed to load: file:///C:/Users/em/Desktop/Surgical%20Relevance%20of%20Corona%20Mortis%20and%20Clinical%20Outcome%20in%20Pelvic%20Trauma-Dateier

Approval by the local ethics committee for database analysis was obtained: "Retrospektive Analysen in der Chirurgischen Intensivmedizin" Nr. St.V 01/2008.

For this retrospective data base study, we enrolled patients, who underwent pelvic surgery due to an acetabular fracture or pelvic ring instability. All patients were treated at our institution. Inclusion criteria were an acetabular fracture and/or pelvic ring instability, senior author participation in the operation and surgery between March 2006 and December 2013. Patients with an age under 15 years, or a pathologic fracture were excluded. In order to further stratify the patients for the evaluation of the CM, any posterior, Olerud and percutaneous surgical approaches were neglected. Up to three approaches were used. The fractures were classified using the classification system described by Young and Burgess [13]. The incidence of the CM was defined by documented intraoperative findings in the operative report. Complications were evaluated based on retrospective chart review of patients' medical records including follow-up radiographs. The minimal follow up time was set at six months. Outcome parameters were the incidence of the CM and its association with bleeding complications, general postoperative complications, and mortality.

## Statistical analysis

All data were recorded in an Excel database (Microsoft Corp., Washington, DC, USA) and statistically analysed with IBM SPSS Statistics for Windows (Version 22.0; IBM Corp., Armonk, NY, USA). Descriptive data are given as mean values and standard deviations or absolute numbers. Normal distribution of the data was tested with a two-tailed Kolmogorov-Smirnov test. Because of the nonparametric distribution, a Mann-Whitney U test was used to test nominal variables. P-values of <0.05 were considered as statistically significant.

## Results

A total of 195 patients were evaluated in this study. The mean age of was  $51 \pm 19$  years and there were 66 (33.8%) females as well as 129 (66.2%) males. The diagnoses and approaches (n=318; 1.6 approaches per patient) are presented in Tables 1-3. In 48.7% (n=95), an anterior approach was used and in 10.8% (n=21), a posterior approach was used. A combination of both approaches was performed in 17.9% (n=35) and an individual approach in 1.5% (n=3), while 29.4% (n=58) were complemented by a percutaneous approach. According to the intraoperative approach and potential documentation of the CM, 130 patients were eventually available for further stratification. The incidence of the CM was 41.5% (n=54). Bleeding complications occurred in 5.1% (n=10) of all patients and 6.2% (n=8) of patients with the CM. The complications are summarized in Table 4. The overall mortality was 8.2% (n=16) and the mortality of patients with the CM was 9.2% (n=12). There was no significant association between the presence of the CM and bleeding complications ( $p=0.068$ ). Furthermore, the presence of the CM did not have an influence on the mortality ( $p=0.068$ ). In accordance with these results, the mortality was not significantly affected by the presence of the CM or bleeding complications ( $p=0.338/0.113$ ).



**Table 1:** Diagnosis (fracture) classified using the classification system described by Young and Burgess of all included cases (n1) and cases with approaches with possible visibility of the CM (n2).



**Table 2:** Detailed description of the approaches of all included cases. Combinations are possible.



**Table 3:** Summarized description of the approaches of all included cases.



**Table 4:** Complications of all included cases (n1=195) and cases with approaches where the CM is potentially visible (n2=130).

## Discussion

The intend of this study was to determine the incidence and the clinical relevance of the CM based on intraoperative findings because previous studies are based on CT results, angiographic findings, and cadaver dissections. Our data show that the CM was documented in 41.5% of the cases, which is consistent with the current literature; although even higher incidences of vascular anastomoses have been reported [1-4]. Against our assumption and despite a bleeding complication rate of 6.2%, the presence of the CM was not associated with increased bleeding complications or higher mortality. Therefore, relevant bleeding might arise not only from the CM, but from other sources such as the posterior part of the pelvis. Another reason might be an ongoing bleeding from cancellous bone exposed from the fracture [14]. In addition, the standard surgical procedure for acetabular fractures and pelvic ring instabilities at our institution includes delayed definite, open surgery with prior external stabilization if necessary, in order to allow auto-tamponade to control the bleeding.

data suggest that there is no need for pre-operative identification if a gentle surgical technique with careful dissection is performed while keeping the high incidence of the CM in mind. This is backed up by the fact that elaborated imaging modalities are timeconsuming modalities, which are often challenging to perform in hemodynamically unstable patients with an urgent need for surgical stabilization [16].

In the present study, we used the classification described by Young and Burgess [13] because it has been shown to have a higher inter- and intraobserver reliability [17,18] than the classification system introduced by Tile [19].

We are aware of several limitations of this study. First, it is a retrospective database study with a rather small sample size. Furthermore, a known limitation for studies dealing with pelvic ring instabilities is that concomitant injuries might influence complications. However, the fact that patients were treated by only one senior surgeon is a clear advantage because of the thorough documentation regarding the presence of the CM. Therefore, the data concerning the incidence of the CM seems to be valid, but data about the outcome should be interpreted with care.

The incidence of the CM is comparable to the current literature, but the clinical relevance in the form of bleeding complications and mortality does not seem to be as high as initially thought.

## Conclusion

The high incidence of the CM was not associated with increased bleeding complications or a higher rate of mortality. Major bleeding in pelvic trauma surgeries may occur at other sites such as the central vessels or venous plexus at the back of the pelvis. Nevertheless, the CM seems to be an important but manageable anatomic structure if handled cautiously with the high incidence in mind; preferably during delayed, definite open surgeries for pelvic trauma.

## References

1. [Smith JC, Gregorius JC, Breazeale BH, Watkins GE. The corona mortis, a frequent vascular variant susceptible to blunt pelvic trauma: identification at routine multidetector CT. Journal of vascular and interventional radiology : JVIR. 2009; 20: 455-460.](#)
2. [Darmanis S, Lewis A, Mansoor A, Bircher M. Corona mortis: an anatomical study with clinical implications in approaches to the pelvis and acetabulum. Clinical anatomy \(New York, NY\). 2007; 20: 433-439.](#)
3. [Berberoglu M, Uz A, Ozmen MM, Bozkurt MC, Erkan C, Taner S, et al. Corona mortis: an anatomic study in seven cadavers and an endoscopic study in 28 patients. Surgical endoscopy. 2001; 15: 72-75.](#)
4. [Drewes PG, Marinis SI, Schaffer JI, Boreham MK, Corton MM. Vascular anatomy over the superior pubic rami in female cadavers. American journal of obstetrics and gynecology. 2005; 193: 2165-2168.](#)
5. [Matta JM. Operative treatment of acetabular fractures through the ilioinguinal approach: a 10-year perspective. Journal of orthopaedic trauma. 2006; 20: S20-S29.](#)
6. [Letournel E. The treatment of acetabular fractures through the ilioinguinal approach. Clinical orthopaedics and related research. 1993; 62-76.](#)
7. [Kloen P, Siebenrock KA, Ganz R. Modification of the ilioinguinal approach. Journal of orthopaedic trauma. 2002; 16: 586-593.](#)
8. [Sagi HC, Afsari A, Dziadosz D. The anterior intra-pelvic \(modified rives-stopppa\) approach for fixation of acetabular fractures. Journal of orthopaedic trauma. 2010; 24: 263-270.](#)
9. [Sarikcioglu L, Sindel M, Akyildiz F, Gur S. Anastomotic vessels in the retropubic region: corona mortis. Folia morphologica. 2003; 62:179-182.](#)
10. [Teague DC, Graney DO, Routt ML, Jr. Retropubic vascular hazards of the ilioinguinal exposure: a cadaveric and clinical study. Journal of orthopaedic trauma. 1996; 10: 156-159.](#)
11. [Hong HX, Pan ZJ, Chen X, Huang ZJ. An anatomical study of corona mortis and its clinical significance. Chinese journal of traumatology = Zhonghua chuang shang za zhi / Chinese Medical Association. 2004; 7: 165-169.](#)
12. [Marsman JW, Schilstra SH, van Leeuwen H. Angiography and embolization of the corona mortis \(aberrant obturator artery\). A source of persistent pelvic bleeding. RoFo : Fortschritte auf dem Gebiete der Rontgenstrahlen und der Nuklearmedizin. 1984; 141: 708-710.](#)
13. [Young JW, Burgess AR, Brumback RJ, Poka A. Pelvic fractures: value of plain radiography in early assessment and management. Radiology. 1986; 160: 445-451.](#)
14. [Magnussen RA, Tressler MA, Obrebsky WT, Kregor PJ. Predicting blood loss in isolated pelvic and acetabular high-energy trauma. Journal of orthopaedic trauma. 2007; 21:603-607.](#)
15. [Karakurt L, Karaca I, Yilmaz E, Burma O, Serin E. Corona mortis: incidence and location. Archives of orthopaedic and trauma surgery. 2002; 122: 163-164.](#)
16. [Pohlemann T, Tosounidis G, Bircher M, Giannoudis P, Culemann U. The German Multicentre Pelvis Registry: a template for an European Expert Network? Injury. 2007; 38: 416-423.](#)
17. [Koo H, Leveridge M, Thompson C, Zdero R, Bhandari M, Kreder HJ, et al. Interobserver reliability of the young-burgess and tile classification systems for fractures of the pelvic ring. Journal of orthopaedic trauma. 2008; 22: 379-384.](#)
18. [Furey AJ, O'Toole RV, Nascone JW, Sciadini MF, Copeland CE, Turen C. Classification of pelvic fractures: analysis of inter- and intraobserver variability using the Young-Burgess and Tile classification systems. Orthopedics. 2009; 32: 401.](#)
19. [Tile M. Pelvic ring fractures: should they be fixed? The Journal of bone and joint surgery British volume. 1988; 70: 1-12.](#)

**Citation:** Jensen KO, Sprengel K, Mica L, Somlyay L, Jentzsch T and Werner CML. Surgical Relevance of Corona Mortis and Clinical Outcome in Pelvic Trauma. Austin J Anat. 2015;2(2): 1033. ISSN : 2381-8921

[Home](#)

[Journal Scope](#)

[Online First](#)

[Current Issue](#)

[Editorial Board](#)

[Instruction for Authors](#)

[Submit Your Article](#)

[Contact Us](#)